Have person lie down

Raise feet 10 to 12 inches.

If he is having trouble breathing raise his head.

Never leave him alone.

Talk to him to assure him that everything will be okay.

Have the person blink.

Pull the upper lid down over the lower one.

Place your thumb on the skin just below the lid and pull it down gently.

Lift the object out with the corner of a sterile gauze pad.

Cover both eyes with a gauze pad and get the patient under medical care.

Bite of Animal

Call police to come pick up the animal.

Bite of Animal

Scrub the bite with plenty of soap and water.

Bite of Animal

Cover the wound with a sterile bandage and get the victim to a doctor.

Take out any foreign matter.

Squeeze gently around the wound.

Wash gently with soap and water.

Apply sterile bandage.

Cut the line and allow a doctor to remove the fishhook.

Push the barb out through the skin.

Snip off the barb with a pair of pliers.

Back the fishhook out of the wound.

Do not break the blisters.

Place the injury in cool water until the pain lessons.

Apply a moist dressing.

Bandage loosely.

Do not apply creams, ointments or sprays.

Symptom: Pale skin.

Symptom: Shallow breathing.

Symptom: Nausea.

Symptom: Dilated pupils, headache and dizziness.

Move patient to a cool shady spot.

Place him on his back and raise his feet.

Fan him and apply cool wet cloths. Give him sips of water.

Poison ivy has three leaves.

Poison oak has three leaves.

Poison sumac has white berries.

Do not scrub.

Rinse with water immediately.

Use Calamine lotion to relieve itching.

Remove lingering sap by laundering clothing.